

DISCRETE SCAN STATISTICS WITH WINDOWS OF ARBITRARY SHAPE

Alexandru Amărioarei

National Institute of Research and Development for Biological Sciences
Bucharest, Romania

MØDAL TEAM - INRIA Lille Nord Europe
Lille, France

The 8th International Workshop on Applied Probability
20-23 June, 2016, Toronto, Canada



OUTLINE

1 INTRODUCTION

- Framework
- Problem

2 METHODOLOGY

- Approximation
- Simulation methods: Normal data

3 SIMULATION STUDY

- Numerical examples
- Power

4 SCANNING THE SURFACE OF A CYLINDER

- Problem
- Approximation
- Numerical Results

5 REFERENCES

OUTLINE

1 INTRODUCTION

- Framework
- Problem

2 METHODOLOGY

- Approximation
- Simulation methods: Normal data

3 SIMULATION STUDY

- Numerical examples
- Power

4 SCANNING THE SURFACE OF A CYLINDER


- Problem
- Approximation
- Numerical Results

5 REFERENCES

Definitions and notations

PRELIMINARY NOTATIONS

Let T_1, T_2 be positive integers



- Rectangular region
 $\mathcal{R}_2 = [0, T_1] \times [0, T_2]$
- $(X_{s_1, s_2})_{\substack{1 \leq s_1 \leq T_1 \\ 1 \leq s_2 \leq T_2}}$ i.i.d. r.v.'s
 - Bernoulli($\mathcal{B}(1, p)$)
 - Binomial($\mathcal{B}(n, p)$)
 - Poisson($\mathcal{P}(\lambda)$)
 - Normal($\mathcal{N}(\mu, \sigma^2)$)
- X_{s_1, s_2} number of observed events
in the elementary subregion
 $r_{s_1, s_2} = [s_1 - 1, s_1] \times [s_2 - 1, s_2]$

TWO DIMENSIONAL SCAN STATISTIC

Let $2 \leq m_s \leq T_s$, $s \in \{1, 2\}$ be positive integers

- Define for $1 \leq i_s \leq T_s - m_s + 1$ and $1 \leq j_s \leq m_s$ the 2-way tensor $\mathcal{X}_{i_1, i_2} \in \mathbb{R}^{m_1 \times m_2}$,

$$\mathcal{X}_{i_1, i_2}(j_1, j_2) = X_{i_1 + j_1 - 1, i_2 + j_2 - 1}$$

- Take $\mathcal{S} : \mathbb{R}^{m_1 \times m_2} \rightarrow \mathbb{R}$ to be a measurable real valued function (*score function*) and define

$$Y_{i_1, i_2}(\mathcal{S}) = \mathcal{S}(\mathcal{X}_{i_1, i_2})$$

DEFINITION



The two dimensional scan statistic with score function \mathcal{S} is defined by

$$S_{m_1, m_2}(T_1, T_2; \mathcal{S}) = \max_{\substack{1 \leq i_1 \leq T_1 - m_1 + 1 \\ 1 \leq i_2 \leq T_2 - m_2 + 1}} Y_{i_1, i_2}(\mathcal{S})$$

SHAPE OF THE SCANNING WINDOW

Let G be the geometrical shape of the scanning window (rectangle, quadrilateral, ellipse, etc.) and \tilde{G} be its corresponding discrete form.


- Rasterization algorithms (omputer vision): continuous shape → discrete shape
 - Line - Bresenham line algorithm ([Bresenham, 1965])
 - Circle - Bresenham circle algorithm ([Bresenham, 1977])
 - Bezier curves - [Foley, 1995]



SHAPE OF THE SCANNING WINDOW

To each discrete shape \tilde{G} it corresponds an unique matrix (2-way tensor) $A(G) = A(\tilde{G})$ (of smallest size) with 0 and 1 entries (1 if there is a point and 0 otherwise):

\tilde{G} : Circle




$A(\tilde{G})$: Circle

			1	1	1	1	1		
			1	1	1	1	1	1	1
			1	1	1	1	1	1	1
			1	1	1	1	1	1	1
			1	1	1	1	1	1	1
			1	1	1	1	1	1	1
			1	1	1	1	1	1	1
			1	1	1	1	1	1	1
			1	1	1	1	1	1	1
			1	1	1	1	1	1	1
			1	1	1	1	1	1	1
			1	1	1	1	1	1	1
			1	1	1	1	1	1	1
			1	1	1	1	1	1	1
			1	1	1	1	1	1	1
			1	1	1	1	1	1	1
			1	1	1	1	1	1	1

SHAPE OF THE SCANNING WINDOW

To each discrete shape \tilde{G} it corresponds an unique matrix (2-way tensor) $A(G) = A(\tilde{G})$ (of smallest size) with 0 and 1 entries (1 if there is a point and 0 otherwise):

$\tilde{G} : \text{Annulus}$



$A(\tilde{G}) : \text{Annulus}$

		1	1	1	1	1			
	1	1	1	1	1	1	1	1	
1	1	1	1	1	1	1	1	1	
1	1	1	1	1	1	1	1	1	
1	1	1	1	1	1	1	1	1	
1	1	1	1	1	1	1	1	1	
1	1	1	1	1	1	1	1	1	
1	1	1	1	1	1	1	1	1	
1	1	1	1	1	1	1	1	1	
1	1	1	1	1	1	1	1	1	

ARBITRARY WINDOW SCAN STATISTIC

Let G be a geometric shape and $A = A(G)$ its corresponding $\{0, 1\}$ matrix of size $m_1 \times m_2$.

- Define the score function \mathcal{S} associated to the shape G by

$$\mathcal{S}(\mathcal{X}_{i_1, i_2}) = A \circ \mathcal{X}_{i_1, i_2} = \sum_{s_1=i_1}^{i_1+m_1-1} \sum_{s_2=i_2}^{i_2+m_2-1} A(s_1 - i_1 + 1, s_2 - i_2 + 1) X_{s_1, s_2}$$

REMARK

If, in particular, the shape G is a rectangle of size $m_1 \times m_2$ than its corresponding $\{0, 1\}$ matrix of the same size has all the entries equal to 1 so the score function

$$\mathcal{S}(\mathcal{X}_{i_1, i_2}) = \sum_{s_1=i_1}^{i_1+m_1-1} \sum_{s_2=i_2}^{i_2+m_2-1} X_{s_1, s_2}$$

is the *classical* rectangular window of the two dimensional scan statistics.

OUTLINE

1 INTRODUCTION

- Framework
- Problem

2 METHODOLOGY

- Approximation
- Simulation methods: Normal data

3 SIMULATION STUDY

- Numerical examples
- Power

4 SCANNING THE SURFACE OF A CYLINDER

- Problem
- Approximation
- Numerical Results

5 REFERENCES

Problem and related work

OBJECTIVE

Find a good estimate for the distribution of the two dimensional discrete scan statistic with score function \mathcal{S}

$$Q_{\mathbf{m}}(\mathbf{T}; \mathcal{S}) = \mathbb{P}(S_{\mathbf{m}}(\mathbf{T}; \mathcal{S}) \leq \tau)$$

with $\mathbf{m} = (m_1, m_2)$ and $\mathbf{T} = (T_1, T_2)$

Previous work:

- Continuous scan statistics
 - Rectangles: [Loader, 1991], [Glaz et al., 2001], [Glaz et al., 2009]
 - Circles: [Anderson and Titterington, 1997]
 - Triangles, ellipses and other convex shapes:
[Alm, 1983, Alm, 1997, Alm, 1998], [Tango and Takahashi, 2005],
[Assunção et al., 2006]
- Discrete scan statistics
 - **No results !**

OUTLINE

1 INTRODUCTION

- Framework
- Problem

2 METHODOLOGY

- Approximation
- Simulation methods: Normal data

3 SIMULATION STUDY

- Numerical examples
- Power

4 SCANNING THE SURFACE OF A CYLINDER

- Problem
- Approximation
- Numerical Results

5 REFERENCES

Approximation methodology for the general scan statistic

APPROXIMATION AND ERROR BOUNDS

THEOREM (GENERALIZATION OF [AMĂRIOAREI, 2014])

Let $t_1, t_2 \in \{2, 3\}$, $Q_{t_1, t_2} = \mathbb{P}(S_m(t_1(m_1 - 1), t_2(m_2 - 1); \mathcal{S}) \leq \tau)$ and $L_s = \left\lfloor \frac{T_s}{m_s - 1} \right\rfloor$, $s \in \{1, 2\}$. If \hat{Q}_{t_1, t_2} is an estimate of Q_{t_1, t_2} , $|\hat{Q}_{t_1, t_2} - Q_{t_1, t_2}| \leq \beta_{t_1, t_2}$ and τ is such that $1 - \hat{Q}_{2,2}(\tau) \leq 0.1$ then

$$\left| \mathbb{P}(S_m(\mathbf{T}; \mathcal{S}) \leq \tau) - \left(2\hat{Q}_2 - \hat{Q}_3 \right) \left[1 + \hat{Q}_2 - \hat{Q}_3 + 2(\hat{Q}_2 - \hat{Q}_3)^2 \right]^{1-L_1} \right| \leq E_{sf} + E_{sapp},$$

where, for $t \in \{2, 3\}$

$$\hat{Q}_t = \left(2\hat{Q}_{t,2} - \hat{Q}_{t,3} \right) \left[1 + \hat{Q}_{t,2} - \hat{Q}_{t,3} + 2(\hat{Q}_{t,2} - \hat{Q}_{t,3})^2 \right]^{1-L_2}$$


$$E_{sf} = (L_1 - 1)(L_2 - 1)(\beta_{2,2} + \beta_{2,3} + \beta_{3,2} + \beta_{3,3})$$

$$E_{sapp} = (L_1 - 1) \left[F_1 \left(1 - \hat{Q}_2 + A_2 + C_2 \right)^2 + (L_2 - 1)(F_2 C_2 + F_3 C_3) \right]$$

$$A_2 = (L_2 - 1)(\beta_{2,2} + \beta_{2,3})$$

$$C_t = (L_2 - 1)F_t \left(1 - \hat{Q}_{t,2} + \beta_{t,2} \right)^2.$$

ILLUSTRATION OF THE APPROXIMATION PROCESS



Find
Approximation

ILLUSTRATION OF THE APPROXIMATION PROCESS





ILLUSTRATION OF THE APPROXIMATION PROCESS



OUTLINE

1 INTRODUCTION

- Framework
- Problem

2 METHODOLOGY

- Approximation
- Simulation methods: Normal data

3 SIMULATION STUDY

- Numerical examples
- Power

4 SCANNING THE SURFACE OF A CYLINDER

- Problem
- Approximation
- Numerical Results

5 REFERENCES

Simulation methods for Normal data

IMPORTANCE SAMPLING ALGORITHM

TEST THE NULL HYPOTHESIS OF RANDOMNESS AGAINST AN ALTERNATIVE OF CLUSTERING

H_0 : The r.v.'s X_{s_1, s_2} are i.i.d. $\mathcal{N}(\mu, \sigma^2)$

H_1 : There exists $\mathcal{R}(i_1, i_2) = [i_1 - 1, i_1 + m_1 - 1] \times [i_2 - 1, i_2 + m_2 - 1] \subset \mathcal{R}_2$ where the r.v.'s $X_{s_1, s_2} \sim \mathcal{N}(\mu_1, \sigma^2)$, $\mu_1 > \mu$ and $X_{s_1, s_2} \sim \mathcal{N}(\mu, \sigma^2)$ outside $\mathcal{R}(i_1, i_2)$

OBJECTIVE

Find a good estimate for $\mathbb{P}_{H_0}(S_m(\mathbf{T}; \mathcal{S}) \geq \tau)$.

We are interested in evaluating the probability

$$\mathbb{P}_{H_0}(S_m(\mathbf{T}; \mathcal{S}) \geq \tau) = \mathbb{P}\left(\bigcup_{i_1=1}^{T_1-m_1+1} \bigcup_{i_2=1}^{T_2-m_2+1} E_{i_1, i_2}(\mathcal{S})\right)$$

where $E_{i_1, i_2}(\mathcal{S}) = \{Y_{i_1, i_2}(\mathcal{S}) \geq \tau\}$.

IMPORTANCE SAMPLING ALGORITHM

Algorithm 1 Importance Sampling Algorithm for Scan Statistics

Begin

Repeat for each k from 1 to $ITER$ (iterations number)

- 1: Generate uniformly the couple $(i_1^{(k)}, i_2^{(k)})$ from the set $\{1, \dots, T_1 - m_1 + 1\} \times \{1, \dots, T_2 - m_2 + 1\}$.
- 2: Given the couple $(i_1^{(k)}, i_2^{(k)})$, generate a sample of the random field $\tilde{\mathbf{X}}^{(k)} = \{\tilde{X}_{s_1, s_2}^{(k)}\}$, with $s_j \in \{1, \dots, T_j\}$ and $j \in \{1, 2\}$, from the conditional distribution of \mathbf{X} given $\left\{Y_{i_1^{(k)}, i_2^{(k)}}(\mathcal{S}) \geq \tau\right\}$.
- 3: Take $c_k = C(\tilde{\mathbf{X}}^{(k)})$ the number of all couples (i_1, i_2) for which $\tilde{Y}_{i_1, i_2}(\mathcal{S}) \geq \tau$ and put $\hat{\rho}_k(2) = \frac{1}{c_k}$.

End Repeat

Return

$$\hat{\rho}(2) = \frac{1}{ITER} \sum_{k=1}^{ITER} \hat{\rho}_k(2), \quad Var[\hat{\rho}(2)] \approx \frac{1}{ITER-1} \sum_{k=1}^{ITER} \left(\hat{\rho}_k(2) - \frac{1}{ITER} \sum_{k=1}^{ITER} \hat{\rho}_k(2) \right)^2$$

End

IMPORTANCE SAMPLING ALGORITHM: $\mathcal{N}(\mu, \sigma^2)$

Step 2 requires to sample:

- $Y_{i_1^{(k)}, i_2^{(k)}}(\mathcal{S})$ from the tail distribution $\mathbb{P}\left(Y_{i_1^{(k)}, i_2^{(k)}}(\mathcal{S}) \geq \tau\right)$ ([Devroye, 1986])
- for the other indices, from the conditional distribution given $\left\{Y_{i_1^{(k)}, i_2^{(k)}}(\mathcal{S}) \geq \tau\right\}$

LEMMA (GENERALIZATION OF [AMĂRIOAREI, 2014, LEMMA 3.4.4])

Let N be a positive integer, $\mathbf{X} = (X_1, X_2, \dots, X_N)$ be a vector of i.i.d. $\mathcal{N}(\mu, \sigma^2)$ and $\mathbf{a} = (a_1, \dots, a_N) \in \mathbb{R}^N$ a non zero constant vector ($a_j \neq 0$ for some particular j). Then conditionally given $\langle \mathbf{a}, \mathbf{X} \rangle = t$, the r.v.'s X_s with $s \neq j$ are jointly distributed as

$$\tilde{X}_s = \frac{a_s}{\|\mathbf{a}\|} \left[\frac{t - \mu a_j}{\|\mathbf{a}\|} - \frac{1}{\|\mathbf{a}\| - |a_j|} \sum_{i \neq j} a_i \left(Z_i - \frac{\mu |a_j|}{\|\mathbf{a}\|} \right) \right] + Z_s$$

where Z_s are i.i.d. $\mathcal{N}(\mu, \sigma^2)$ r.v.s.

OUTLINE

1 INTRODUCTION

- Framework
- Problem

2 METHODOLOGY

- Approximation
- Simulation methods: Normal data

3 SIMULATION STUDY


- Numerical examples
- Power

4 SCANNING THE SURFACE OF A CYLINDER

- Problem
- Approximation
- Numerical Results

5 REFERENCES


Numerical examples

SCANNING A REGION OF SIZE $T_1 \times T_2 = 250 \times 250$ TABLE 1 : Numerical results for $\mathbb{P}(S \leq \tau)$: Triangle


Window's shape		Triangle ($m_1 = 14, m_2 = 18, Nt = 133, IS = 1e5, IA = 1e6$)							
		$X_{s_1, s_2} \sim \mathcal{B}(1, 0.01)$			$X_{s_1, s_2} \sim \mathcal{B}(5, 0.05)$				
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal		
3	0.916397	0.918667	0.004333	59	0.863336	0.897101	0.004902		
4	0.997488	0.997483	0.000082	60	0.936684	0.948160	0.002010		
5	0.999947	0.999947	0.000002	61	0.971529	0.974938	0.000894		
6	0.999999	0.999999	0	62	0.987094	0.988279	0.000412		
7	0.999999	0.999999	0	63	0.994372	0.994664	0.000192		
8	1.000000	1.000000	0	64	0.997563	0.997643	0.000089		
9	1.000000	1.000000	0	65	0.998946	0.998971	0.000041		
10	1.000000	1.000000	0	66	0.999564	0.999567	0.000018		
11	1.000000	1.000000	0	67	0.999817	0.999820	0.000008		
		$X_{s_1, s_2} \sim \mathcal{P}(0.25)$			$X_{s_1, s_2} \sim \mathcal{N}(0, 1)$				
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal		
59	0.739866	0.820277	0.011813	50	0.911170	0.934999	0.002737		
60	0.871829	0.902957	0.004483	51	0.945219	0.957777	0.001655		
61	0.939577	0.950977	0.001911	52	0.966494	0.972997	0.001026		
62	0.971673	0.975649	0.000890	53	0.979944	0.982929	0.000644		
63	0.987075	0.988206	0.000425	54	0.987840	0.989469	0.000406		
64	0.994104	0.994492	0.000204	55	0.992768	0.993477	0.000257		
65	0.997384	0.997452	0.000098	56	0.995667	0.996022	0.000162		
66	0.998821	0.998855	0.000046	57	0.997412	0.997574	0.000102		
67	0.999489	0.999490	0.000022	58	0.998509	0.998563	0.000063		

SCANNING A REGION OF SIZE $T_1 \times T_2 = 250 \times 250$ TABLE 1 : Numerical results for $\mathbb{P}(S \leq \tau)$: Triangle

Window's shape


Triangle ($m_1 = 14, m_2 = 18, Nt = 133, IS = 1e5, IA = 1e6$)

$X_{s_1, s_2} \sim \mathcal{B}(1, 0.01)$				$X_{s_1, s_2} \sim \mathcal{B}(5, 0.05)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal
3	0.916397	0.918667	0.004333	59	0.863336	0.897101	0.004902
4	0.997488	0.997483	0.000082	60	0.936684	0.948160	0.002010
5	0.999947	0.999947	0.000002	61	0.971529	0.974938	0.000894
6	0.999999	0.999999	0	62	0.987094	0.988279	0.000412
7	0.999999	0.999999	0	63	0.994372	0.994664	0.000192
8	1.000000	1.000000	0	64	0.997563	0.997643	0.000089
9	1.000000	1.000000	0	65	0.998946	0.998971	0.000041
10	1.000000	1.000000	0	66	0.999564	0.999567	0.000018
11	1.000000	1.000000	0	67	0.999817	0.999820	0.000008


$X_{s_1, s_2} \sim \mathcal{P}(0.25)$				$X_{s_1, s_2} \sim \mathcal{N}(0, 1)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal
59	0.739866	0.820277	0.011813	50	0.911170	0.934999	0.002737
60	0.871829	0.902957	0.004483	51	0.945219	0.957777	0.001655
61	0.939577	0.950977	0.001911	52	0.966494	0.972997	0.001026
62	0.971673	0.975649	0.000890	53	0.979944	0.982929	0.000644
63	0.987075	0.988206	0.000425	54	0.987840	0.989469	0.000406
64	0.994104	0.994492	0.000204	55	0.992768	0.993477	0.000257
65	0.997384	0.997452	0.000098	56	0.995667	0.996022	0.000162
66	0.998821	0.998855	0.000046	57	0.997412	0.997574	0.000102
67	0.999489	0.999490	0.000022	58	0.9998509	0.9998563	0.000063



SCANNING A REGION OF SIZE $T_1 \times T_2 = 250 \times 250$ TABLE 2 : Numerical results for $\mathbb{P}(S \leq \tau)$: Rectangle

Window's shape		Rectangle ($m_1 = 11, m_2 = 12, Nt = 132, IS = 1e5, IA = 1e6$)						
		$X_{s_1, s_2} \sim \mathcal{B}(1, 0.01)$			$X_{s_1, s_2} \sim \mathcal{B}(5, 0.05)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal	
	3	0.947896	0.948226	0.002110	59	0.856993	0.857094	0.005485
	4	0.997983	0.997986	0.000058	60	0.919416	0.919589	0.002300
	5	0.999943	0.999943	0.000001	61	0.956420	0.956569	0.001024
	6	0.999999	0.999999	0	62	0.977142	0.977065	0.000471
	7	0.999999	0.999999	0	63	0.988228	0.988208	0.000220
	8	1.000000	1.000000	0	64	0.994130	0.994095	0.000103
	9	1.000000	1.000000	0	65	0.997121	0.997107	0.000048
	10	1.000000	1.000000	0	66	0.998610	0.998607	0.000022
	11	1.000000	1.000000	0	67	0.999342	0.999342	0.000010
		$X_{s_1, s_2} \sim \mathcal{P}(0.25)$			$X_{s_1, s_2} \sim \mathcal{N}(0, 1)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal	
59	0.764372	0.764495	0.013285	50	0.865858	0.865484	0.004572	
60	0.857872	0.859289	0.005320	51	0.904555	0.904805	0.002691	
61	0.918972	0.918732	0.002307	52	0.933323	0.933206	0.001620	
62	0.954682	0.954579	0.001059	53	0.953950	0.953807	0.000993	
63	0.975271	0.975391	0.000502	54	0.968340	0.968483	0.000617	
64	0.986996	0.986966	0.000242	55	0.978632	0.978687	0.000386	
65	0.993240	0.993261	0.000117	56	0.985791	0.985752	0.000242	
66	0.996557	0.996551	0.000056	57	0.990621	0.990585	0.000152	
67	0.998290	0.998283	0.000027	58	0.993837	0.993815	0.000096	

SCANNING A REGION OF SIZE $T_1 \times T_2 = 250 \times 250$ TABLE 2 : Numerical results for $\mathbb{P}(S \leq \tau)$: Rectangle

Window's shape		Rectangle ($m_1 = 11, m_2 = 12, Nt = 132, IS = 1e5, IA = 1e6$)						
		$X_{s_1, s_2} \sim \mathcal{B}(1, 0.01)$			$X_{s_1, s_2} \sim \mathcal{B}(5, 0.05)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal	
	3	0.947896	0.948226	0.002110	59	0.856993	0.857094	0.005485
	4	0.997983	0.997986	0.000058	60	0.919416	0.919589	0.002300
	5	0.999943	0.999943	0.000001	61	0.956420	0.956569	0.001024
	6	0.999999	0.999999	0	62	0.977142	0.977065	0.000471
	7	0.999999	0.999999	0	63	0.988228	0.988208	0.000220
	8	1.000000	1.000000	0	64	0.994130	0.994095	0.000103
	9	1.000000	1.000000	0	65	0.997121	0.997107	0.000048
	10	1.000000	1.000000	0	66	0.998610	0.998607	0.000022
	11	1.000000	1.000000	0	67	0.999342	0.999342	0.000010
		$X_{s_1, s_2} \sim \mathcal{P}(0.25)$			$X_{s_1, s_2} \sim \mathcal{N}(0, 1)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal	
59	0.764372	0.764495	0.013285	50	0.865858	0.865484	0.004572	
60	0.857872	0.859289	0.005320	51	0.904555	0.904805	0.002691	
61	0.918972	0.918732	0.002307	52	0.933323	0.933206	0.001620	
62	0.954682	0.954579	0.001059	53	0.953950	0.953807	0.000993	
63	0.975271	0.975391	0.000502	54	0.968340	0.968483	0.000617	
64	0.986996	0.986966	0.000242	55	0.978632	0.978687	0.000386	
65	0.993240	0.993261	0.000117	56	0.985791	0.985752	0.000242	
66	0.996557	0.996551	0.000056	57	0.990621	0.990585	0.000152	
67	0.998290	0.998283	0.000027	58	0.993837	0.993815	0.000096	

SCANNING A REGION OF SIZE $T_1 \times T_2 = 250 \times 250$ TABLE 3 : Numerical results for $\mathbb{P}(S \leq \tau)$: Quadrilateral


Window's shape		Quadrilateral ($m_1 = 14, m_2 = 18, Nt = 131, IS = 1e5, IA = 1e6$)						
		$X_{s_1, s_2} \sim \mathcal{B}(1, 0.01)$			$X_{s_1, s_2} \sim \mathcal{B}(5, 0.05)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal	
3	0.926068	0.927398	0.003806	59	0.914546	0.927613	0.002942	
4	0.997622	0.997627	0.000075	60	0.959599	0.963873	0.001255	
5	0.999946	0.999946	0.000002	61	0.981235	0.982506	0.000571	
6	0.999999	0.999999	0	62	0.991423	0.991796	0.000266	
7	0.999999	0.999999	0	63	0.996113	0.996233	0.000124	
8	1.000000	1.000000	0	64	0.998283	0.998337	0.000057	
9	1.000000	1.000000	0	65	0.999266	0.999266	0.000026	
10	1.000000	1.000000	0	66	0.999684	0.999684	0.000012	
11	1.000000	1.000000	0	67	0.999868	0.999869	0.000005	
		$X_{s_1, s_2} \sim \mathcal{P}(0.25)$			$X_{s_1, s_2} \sim \mathcal{N}(0, 1)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal	
59	0.835054	0.870351	0.006852	50	0.920004	0.935266	0.002571	
60	0.917972	0.931040	0.002768	51	0.950232	0.957711	0.001556	
61	0.960397	0.964711	0.001237	52	0.968755	0.972594	0.000964	
62	0.981228	0.982451	0.000585	53	0.980695	0.982566	0.000606	
63	0.991142	0.991510	0.000281	54	0.988110	0.989060	0.000383	
64	0.995855	0.995971	0.000136	55	0.992626	0.993110	0.000242	
65	0.998108	0.998124	0.000065	56	0.995569	0.995771	0.000153	
66	0.999135	0.999153	0.000031	57	0.997361	0.997394	0.000096	
67	0.999620	0.999622	0.000014	58	0.998379	0.998435	0.000060	

SCANNING A REGION OF SIZE $T_1 \times T_2 = 250 \times 250$ TABLE 3 : Numerical results for $\mathbb{P}(S \leq \tau)$: Quadrilateral

Window's shape		Quadrilateral ($m_1 = 14, m_2 = 18, Nt = 131, IS = 1e5, IA = 1e6$)						
		$X_{s_1, s_2} \sim \mathcal{B}(1, 0.01)$			$X_{s_1, s_2} \sim \mathcal{B}(5, 0.05)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal	
3	0.926068	0.927398	0.003806	59	0.914546	0.927613	0.002942	
4	0.997622	0.997627	0.000075	60	0.959599	0.963873	0.001255	
5	0.999946	0.999946	0.000002	61	0.981235	0.982506	0.000571	
6	0.999999	0.999999	0	62	0.991423	0.991796	0.000266	
7	0.999999	0.999999	0	63	0.996113	0.996233	0.000124	
8	1.000000	1.000000	0	64	0.998283	0.998337	0.000057	
9	1.000000	1.000000	0	65	0.999266	0.999266	0.000026	
10	1.000000	1.000000	0	66	0.999684	0.999684	0.000012	
11	1.000000	1.000000	0	67	0.999868	0.999869	0.000005	
		$X_{s_1, s_2} \sim \mathcal{P}(0.25)$			$X_{s_1, s_2} \sim \mathcal{N}(0, 1)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal	
59	0.835054	0.870351	0.006852	50	0.920004	0.935266	0.002571	
60	0.917972	0.931040	0.002768	51	0.950232	0.957711	0.001556	
61	0.960397	0.964711	0.001237	52	0.968755	0.972594	0.000964	
62	0.981228	0.982451	0.000585	53	0.980695	0.982566	0.000606	
63	0.991142	0.991510	0.000281	54	0.988110	0.989060	0.000383	
64	0.995855	0.995971	0.000136	55	0.992626	0.993110	0.000242	
65	0.998108	0.998124	0.000065	56	0.995569	0.995771	0.000153	
66	0.999135	0.999153	0.000031	57	0.997361	0.997394	0.000096	
67	0.999620	0.999622	0.000014	58	0.998379	0.998435	0.000060	

SCANNING A REGION OF SIZE $T_1 \times T_2 = 250 \times 250$


TABLE 4 : Numerical results for $\mathbb{P}(S \leq \tau)$: Circle

Window's shape		Circle ($m_1 = 13, m_2 = 13, Nt = 129, IS = 1e54, IA = 1e6$)					
		$X_{s_1, s_2} \sim \mathcal{B}(1, 0.01)$			$X_{s_1, s_2} \sim \mathcal{B}(5, 0.05)$		
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal
3	0.950311	0.950461	0.002195	59 60 61 62 63 64 65 66 67	0.920229	0.920388	0.002318
4	0.998118	0.998114	0.000059		0.956814	0.957143	0.001016
5	0.999947	0.999947	0.000001		0.977460	0.977614	0.000462
6	0.999999	0.999999	0		0.988568	0.988567	0.000214
7	0.999999	0.999999	0		0.994312	0.994309	0.000099
8	1.000000	1.000000	0		0.997229	0.997228	0.000046
9	1.000000	1.000000	0		0.998678	0.998679	0.000021
10	1.000000	1.000000	0		0.999380	0.999381	0.000009
11	1.000000	1.000000	0		0.999715	0.999715	0.000004
		$X_{s_1, s_2} \sim \mathcal{P}(0.25)$			$X_{s_1, s_2} \sim \mathcal{N}(0, 1)$		
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal
59	0.858454	0.859178	0.005391	50	0.888137	0.887891	0.003485
60	0.919182	0.919586	0.002310	51	0.921173	0.921549	0.002058
61	0.955229	0.955388	0.001047	52	0.945761	0.945644	0.001243
62	0.976023	0.975987	0.000491	53	0.962790	0.962825	0.000760
63	0.987414	0.987344	0.000234	54	0.974848	0.974878	0.000470
64	0.993477	0.993502	0.000112	55	0.983235	0.983263	0.000293
65	0.996706	0.996703	0.000054	56	0.988885	0.988907	0.000182
66	0.998372	0.998365	0.000025	57	0.992730	0.992734	0.000114
67	0.999207	0.999203	0.000012	58	0.995269	0.995287	0.000071

SCANNING A REGION OF SIZE $T_1 \times T_2 = 250 \times 250$


TABLE 4 : Numerical results for $\mathbb{P}(S \leq \tau)$: Circle

Window's shape		Circle ($m_1 = 13, m_2 = 13, Nt = 129, IS = 1e54, IA = 1e6$)					
		$X_{s_1, s_2} \sim \mathcal{B}(1, 0.01)$			$X_{s_1, s_2} \sim \mathcal{B}(5, 0.05)$		
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal
3	0.950311	0.950461	0.002195	59	0.920229	0.920388	0.002318
4	0.998118	0.998114	0.000059	60	0.956814	0.957143	0.001016
5	0.999947	0.999947	0.000001	61	0.977460	0.977614	0.000462
6	0.999999	0.999999	0	62	0.988568	0.988567	0.000214
7	0.999999	0.999999	0	63	0.994312	0.994309	0.000099
8	1.000000	1.000000	0	64	0.997229	0.997228	0.000046
9	1.000000	1.000000	0	65	0.998678	0.998679	0.000021
10	1.000000	1.000000	0	66	0.999380	0.999381	0.000009
11	1.000000	1.000000	0	67	0.999715	0.999715	0.000004
		$X_{s_1, s_2} \sim \mathcal{P}(0.25)$			$X_{s_1, s_2} \sim \mathcal{N}(0, 1)$		
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal
59	0.858454	0.859178	0.005391	50	0.888137	0.887891	0.003485
60	0.919182	0.919586	0.002310	51	0.921173	0.921549	0.002058
61	0.955229	0.955388	0.001047	52	0.945761	0.945644	0.001243
62	0.976023	0.975987	0.000491	53	0.962790	0.962825	0.000760
63	0.987414	0.987344	0.000234	54	0.974848	0.974878	0.000470
64	0.993477	0.993502	0.000112	55	0.983235	0.983263	0.000293
65	0.996706	0.996703	0.000054	56	0.988885	0.988907	0.000182
66	0.998372	0.998365	0.000025	57	0.992730	0.992734	0.000114
67	0.999207	0.9992032	0.000012	58	0.995269	0.995287	0.000071



SCANNING A REGION OF SIZE $T_1 \times T_2 = 250 \times 250$

TABLE 5 : Numerical results for $\mathbb{P}(S \leq \tau)$: Ellipse

Window's shape		Ellipse ($m_1 = 19, m_2 = 9, Nt = 135, IS = 1e5, IA = 1e6$)						
		$X_{s_1, s_2} \sim \mathcal{B}(1, 0.01)$			$X_{s_1, s_2} \sim \mathcal{B}(5, 0.05)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal	
	3	0.944001	0.944211	0.002297	59	0.764871	0.763482	0.009128
	4	0.997757	0.997758	0.000069	60	0.860903	0.860548	0.004127
	5	0.999935	0.999935	0.000002	61	0.921089	0.920882	0.001941
	6	0.999998	0.999998	0	62	0.956735	0.956866	0.000934
	7	1.000000	1.000000	0	63	0.977118	0.977094	0.000452
	8	1.000000	1.000000	0	64	0.988182	0.988152	0.000218
	9	1.000000	1.000000	0	65	0.994044	0.994012	0.000104
	10	1.000000	1.000000	0	66	0.997037	0.997037	0.000049
	11	1.000000	1.000000	0	67	0.998554	0.998558	0.000023
		$X_{s_1, s_2} \sim \mathcal{P}(0.25)$			$X_{s_1, s_2} \sim \mathcal{N}(0, 1)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal	
59	0.638962	0.639102	0.019442	50	0.843156	0.844113	0.004369	
60	0.768283	0.769666	0.008610	51	0.887477	0.887692	0.002755	
61	0.861614	0.860885	0.004012	52	0.920601	0.920385	0.001757	
62	0.919144	0.919301	0.001948	53	0.944398	0.944328	0.001127	
63	0.954941	0.954864	0.000965	54	0.961682	0.961667	0.000725	
64	0.975255	0.975369	0.000481	55	0.973797	0.973864	0.000468	
65	0.986869	0.986900	0.000240	56	0.982232	0.982330	0.000301	
66	0.993115	0.993127	0.000119	57	0.988183	0.988132	0.000193	
67	0.996485	0.996472	0.000058	58	0.992138	0.992134	0.000123	

SCANNING A REGION OF SIZE $T_1 \times T_2 = 250 \times 250$


TABLE 5 : Numerical results for $\mathbb{P}(S \leq \tau)$: Ellipse

Window's shape		Ellipse ($m_1 = 19, m_2 = 9, Nt = 135, IS = 1e5, IA = 1e6$)					
		$X_{s_1, s_2} \sim \mathcal{B}(1, 0.01)$			$X_{s_1, s_2} \sim \mathcal{B}(5, 0.05)$		
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal
3	0.944001	0.944211	0.002297	59	0.764871	0.763482	0.009128
4	0.997757	0.997758	0.000069	60	0.860903	0.860548	0.004127
5	0.999935	0.999935	0.000002	61	0.921089	0.920882	0.001941
6	0.999998	0.999998	0	62	0.956735	0.956866	0.000934
7	1.000000	1.000000	0	63	0.977118	0.977094	0.000452
8	1.000000	1.000000	0	64	0.988182	0.988152	0.000218
9	1.000000	1.000000	0	65	0.994044	0.994012	0.000104
10	1.000000	1.000000	0	66	0.997037	0.997037	0.000049
11	1.000000	1.000000	0	67	0.998554	0.998558	0.000023
		$X_{s_1, s_2} \sim \mathcal{P}(0.25)$			$X_{s_1, s_2} \sim \mathcal{N}(0, 1)$		
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal
59	0.638962	0.639102	0.019442	50	0.843156	0.844113	0.004369
60	0.768283	0.769666	0.008610	51	0.887477	0.887692	0.002755
61	0.861614	0.860885	0.004012	52	0.920601	0.920385	0.001757
62	0.919144	0.919301	0.001948	53	0.944398	0.944328	0.001127
63	0.954941	0.954864	0.000965	54	0.961682	0.961667	0.000725
64	0.975255	0.975369	0.000481	55	0.973797	0.973864	0.000468
65	0.986869	0.986900	0.000240	56	0.982232	0.982330	0.000301
66	0.993115	0.993127	0.000119	57	0.988183	0.988132	0.000193
67	0.996485	0.996472	0.000058	58	0.992138	0.992134	0.000123

SCANNING A REGION OF SIZE $T_1 \times T_2 = 250 \times 250$


TABLE 6 : Numerical results for $\mathbb{P}(S \leq \tau)$: Ellipse2

Window's shape		Ellipse2 ($m_1 = 9, m_2 = 19, Nt = 135, IS = 1e5, IA = 1e6$)					
		$X_{s_1, s_2} \sim \mathcal{B}(1, 0.01)$			$X_{s_1, s_2} \sim \mathcal{B}(5, 0.05)$		
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal
3	0.944180	0.943779	0.003420	59	0.764090	0.763753	0.036701
4	0.997761	0.997767	0.000071	60	0.859511	0.860602	0.011219
5	0.999934	0.999935	0.000002	61	0.921173	0.920669	0.003816
6	0.999998	0.999998	0	62	0.956920	0.956693	0.001440
7	0.999999	0.999999	0	63	0.977198	0.977129	0.000586
8	1.000000	1.000000	0	64	0.988162	0.988177	0.000253
9	1.000000	1.000000	0	65	0.993979	0.994008	0.000113
10	1.000000	1.000000	0	66	0.997032	0.997036	0.000051
11	1.000000	1.000000	0	67	0.998558	0.998562	0.000023
		$X_{s_1, s_2} \sim \mathcal{P}(0.25)$			$X_{s_1, s_2} \sim \mathcal{N}(0, 1)$		
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal
59	0.638389	0.640805	0.117483	50	0.843792	0.844342	0.013346
60	0.770899	0.769795	0.034030	51	0.888070	0.887823	0.006857
61	0.860847	0.861092	0.010890	52	0.920460	0.920691	0.003626
62	0.919522	0.919537	0.003909	53	0.944514	0.944368	0.001974
63	0.954873	0.954742	0.001516	54	0.961591	0.961748	0.001109
64	0.975326	0.975379	0.000635	55	0.973861	0.973830	0.000640
65	0.986856	0.986843	0.000282	56	0.982294	0.982269	0.000377
66	0.993154	0.993119	0.000130	57	0.988180	0.988143	0.000226
67	0.996482	0.996478	0.000061	58	0.992154	0.992123	0.000138



SCANNING A REGION OF SIZE $T_1 \times T_2 = 250 \times 250$ TABLE 6 : Numerical results for $\mathbb{P}(S \leq \tau)$: Ellipse2

Window's shape

Ellipse2 ($m_1 = 9, m_2 = 19, Nt = 135, IS = 1e5, IA = 1e6$)

$X_{s_1, s_2} \sim \mathcal{B}(1, 0.01)$				$X_{s_1, s_2} \sim \mathcal{B}(5, 0.05)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal
3	0.944180	0.943779	0.003420	59	0.764090	0.763753	0.036701
4	0.997761	0.997767	0.000071	60	0.859511	0.860602	0.011219
5	0.999934	0.999935	0.000002	61	0.921173	0.920669	0.003816
6	0.999998	0.999998	0	62	0.956920	0.956693	0.001440
7	0.999999	0.999999	0	63	0.977198	0.977129	0.000586
8	1.000000	1.000000	0	64	0.988162	0.988177	0.000253
9	1.000000	1.000000	0	65	0.993979	0.994008	0.000113
10	1.000000	1.000000	0	66	0.997032	0.997036	0.000051
11	1.000000	1.000000	0	67	0.998558	0.998562	0.000023

 $X_{s_1, s_2} \sim \mathcal{P}(0.25)$ $X_{s_1, s_2} \sim \mathcal{N}(0, 1)$

τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal
59	0.638389	0.640805	0.117483	50	0.843792	0.844342	0.013346
60	0.770899	0.769795	0.034030	51	0.888070	0.887823	0.006857
61	0.860847	0.861092	0.010890	52	0.920460	0.920691	0.003626
62	0.919522	0.919537	0.003909	53	0.944514	0.944368	0.001974
63	0.954873	0.954742	0.001516	54	0.961591	0.961748	0.001109
64	0.975326	0.975379	0.000635	55	0.973861	0.973830	0.000640
65	0.986856	0.986843	0.000282	56	0.982294	0.982269	0.000377
66	0.993154	0.993119	0.000130	57	0.988180	0.988143	0.000226
67	0.996482	0.996478	0.000061	58	0.992154	0.992123	0.000138

SCANNING A REGION OF SIZE $T_1 \times T_2 = 250 \times 250$

TABLE 7 : Numerical results for $\mathbb{P}(S \leq \tau)$: Annulus

Window's shape		Annulus ($m_1 = 17$, $m_2 = 17$, $Nt = 124$, $IS = 1e5$, $IA = 1e6$)						
		$X_{s_1, s_2} \sim \mathcal{B}(1, 0.01)$			$X_{s_1, s_2} \sim \mathcal{B}(5, 0.05)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal	
	3	0.881798	0.882489	0.004812	59	0.951170	0.951245	0.000699
	4	0.995434	0.995465	0.000069	60	0.975772	0.975727	0.000255
	5	0.999883	0.999883	0.000001	61	0.988275	0.988270	0.000099
	6	0.999998	0.999998	0	62	0.994460	0.994466	0.000041
	7	0.999999	0.999999	0	63	0.997439	0.997440	0.000017
	8	1.000000	1.000000	0	64	0.998839	0.998840	0.000007
	9	1.000000	1.000000	0	65	0.999484	0.999484	0.000003
	10	1.000000	1.000000	0	66	0.999775	0.999775	0.000001
	11	1.000000	1.000000	0	67	0.999903	0.999903	0.000000
		$X_{s_1, s_2} \sim \mathcal{P}(0.25)$			$X_{s_1, s_2} \sim \mathcal{N}(0, 1)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal	
59	0.903956	0.903852	0.002128	50	0.860708	0.860416	0.004097	
60	0.949083	0.949059	0.000735	51	0.904651	0.904644	0.001977	
61	0.973814	0.973844	0.000277	52	0.936077	0.935938	0.000987	
62	0.986909	0.986876	0.000111	53	0.957564	0.957650	0.000508	
63	0.993576	0.993570	0.000047	54	0.972378	0.972311	0.000270	
64	0.996910	0.996907	0.000020	55	0.982139	0.982136	0.000148	
65	0.998539	0.998539	0.000009	56	0.988564	0.988587	0.000082	
66	0.999320	0.999320	0.000004	57	0.992769	0.992769	0.000047	
67	0.999689	0.999689	0.000002	58	0.995471	0.995466	0.000027	

SCANNING A REGION OF SIZE $T_1 \times T_2 = 250 \times 250$ TABLE 7 : Numerical results for $\mathbb{P}(S \leq \tau)$: Annulus

Window's shape		Annulus ($m_1 = 17$, $m_2 = 17$, $Nt = 124$, $IS = 1e5$, $IA = 1e6$)						
		$X_{s_1, s_2} \sim \mathcal{B}(1, 0.01)$			$X_{s_1, s_2} \sim \mathcal{B}(5, 0.05)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal	
	3	0.881798	0.882489	0.004812	59	0.951170	0.951245	0.000699
	4	0.995434	0.995465	0.000069	60	0.975772	0.975727	0.000255
	5	0.999883	0.999883	0.000001	61	0.988275	0.988270	0.000099
	6	0.999998	0.999998	0	62	0.994460	0.994466	0.000041
	7	0.999999	0.999999	0	63	0.997439	0.997440	0.000017
	8	1.000000	1.000000	0	64	0.998839	0.998840	0.000007
	9	1.000000	1.000000	0	65	0.999484	0.999484	0.000003
	10	1.000000	1.000000	0	66	0.999775	0.999775	0.000001
	11	1.000000	1.000000	0	67	0.999903	0.999903	0.000000
		$X_{s_1, s_2} \sim \mathcal{P}(0.25)$			$X_{s_1, s_2} \sim \mathcal{N}(0, 1)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal	
59	0.903956	0.903852	0.002128	50	0.860708	0.860416	0.004097	
60	0.949083	0.949059	0.000735	51	0.904651	0.904644	0.001977	
61	0.973814	0.973844	0.000277	52	0.936077	0.935938	0.000987	
62	0.986909	0.986876	0.000111	53	0.957564	0.957650	0.000508	
63	0.993576	0.993570	0.000047	54	0.972378	0.972311	0.000270	
64	0.996910	0.996907	0.000020	55	0.982139	0.982136	0.000148	
65	0.998539	0.998539	0.000009	56	0.988564	0.988587	0.000082	
66	0.999320	0.999320	0.000004	57	0.992769	0.992769	0.000047	
67	0.999689	0.999689	0.000002	58	0.995471	0.995466	0.000027	

OUTLINE

1 INTRODUCTION

- Framework
- Problem

2 METHODOLOGY

- Approximation
- Simulation methods: Normal data

3 SIMULATION STUDY

- Numerical examples
- Power

4 SCANNING THE SURFACE OF A CYLINDER


- Problem
- Approximation
- Numerical Results

5 REFERENCES


Power of the scan statistic test

POWER EVALUATION FOR $\mathcal{B}(1, 0.001)$ MODEL

Triangular simulated cluster




Rectangular simulated cluster




POWER EVALUATION FOR $\mathcal{B}(1, 0.001)$ MODEL

Quadrilateral simulated cluster




Circular simulated cluster




POWER EVALUATION FOR $\mathcal{B}(1, 0.001)$ MODEL

Ellipsoidal simulated cluster




Annular simulated cluster




POWER EVALUATION FOR $\mathcal{B}(5, 0.05)$ MODEL

Triangular simulated cluster




Rectangular simulated cluster




POWER EVALUATION FOR $\mathcal{B}(5, 0.05)$ MODEL

Quadrilateral simulated cluster




Circular simulated cluster




POWER EVALUATION FOR $\mathcal{B}(5, 0.05)$ MODEL

Ellipsoidal simulated cluster



Annular simulated cluster



OUTLINE

1 INTRODUCTION

- Framework
- Problem

2 METHODOLOGY

- Approximation
- Simulation methods: Normal data

3 SIMULATION STUDY

- Numerical examples
- Power




4 SCANNING THE SURFACE OF A CYLINDER

- Problem
- Approximation
- Numerical Results

5 REFERENCES

Scanning the surface of a cylinder

SCANNING THE SURFACE OF A CYLINDER

Unfolded cylinder of size $T_1 \times T_2$ 

Inria

OUTLINE

1 INTRODUCTION

- Framework
- Problem

2 METHODOLOGY

- Approximation
- Simulation methods: Normal data

3 SIMULATION STUDY

- Numerical examples
- Power


4 SCANNING THE SURFACE OF A CYLINDER

- Problem
- Approximation
- Numerical Results


5 REFERENCES

Transformation into a one dimensional problem


TRANSFORMATION OF THE UNFOLDED CYLINDER




TRANSFORMATION OF THE UNFOLDED CYLINDER




TRANSFORMATION OF THE UNFOLDED CYLINDER




TRANSFORMATION OF THE UNFOLDED CYLINDER




TRANSFORMATION OF THE UNFOLDED CYLINDER



TRANSFORMATION OF THE UNFOLDED CYLINDER




TRANSFORMATION OF THE UNFOLDED CYLINDER



$x_{1,1}$	\dots	$x_{m_1,1}$	\dots	$x_{T_1,1}$	\dots	x_{1,m_2}	\dots	x_{m_1,m_2}	\dots	x_{T_1,m_2}	\dots	\dots	x_{1,T_2}	\dots	x_{m_1,T_2}	\dots	x_{T_1,T_2}
-----------	---------	-------------	---------	-------------	---------	-------------	---------	---------------	---------	---------------	---------	---------	-------------	---------	---------------	---------	---------------

DEFINING THE SCORE FUNCTION




- The size of the scanning window is $\tilde{m}_1 = T_1 m_2 - (T_1 - m_1)$
- The score function is defined by

$$\mathcal{S}(\mathbf{x}_{i_1}) = A \circ \mathbf{x}_{i_1}$$

where A is the corresponding $\{0, 1\}$ vector

DEFINING THE SCORE FUNCTION




- The size of the scanning window is $\tilde{m}_1 = T_1 m_2 - (T_1 - m_1)$
- The score function is defined by

$$\mathcal{S}(\mathbf{x}_{i_1}) = A \circ \mathbf{x}_{i_1}$$

where A is the corresponding $\{0, 1\}$ vector

DEFINING THE SCORE FUNCTION




- The size of the scanning window is $\tilde{m}_1 = T_1 m_2 - (T_1 - m_1)$
- The score function is defined by

$$\mathcal{S}(\mathbf{x}_{i_1}) = A \circ \mathbf{x}_{i_1}$$

where A is the corresponding $\{0, 1\}$ vector



DEFINING THE SCORE FUNCTION




- The size of the scanning window is $\tilde{m}_1 = T_1 m_2 - (T_1 - m_1)$
- The score function is defined by

$$\mathcal{S}(\mathbf{x}_{i_1}) = A \circ \mathbf{x}_{i_1}$$

where A is the corresponding $\{0, 1\}$ vector



DEFINING THE SCORE FUNCTION




- The size of the scanning window is $\tilde{m}_1 = T_1 m_2 - (T_1 - m_1)$
- The score function is defined by

$$\mathcal{S}(\mathbf{x}_{i_1}) = A \circ \mathbf{x}_{i_1}$$

where A is the corresponding $\{0, 1\}$ vector



DEFINING THE SCORE FUNCTION



- The size of the scanning window is $\tilde{m}_1 = T_1 m_2 - (T_1 - m_1)$
- The score function is defined by

$$\mathcal{S}(\mathbf{x}_{i_1}) = A \circ \mathbf{x}_{i_1}$$

where A is the corresponding $\{0, 1\}$ vector



APPROXIMATION AND ERROR BOUNDS


THEOREM [AMĂRIOAREI, 2014]

Let $t_1 \in \{2, 3\}$ and $Q_{t_1} = Q_{t_1}(\tau) = \mathbb{P}(S_{\tilde{m}_1}(t_1(\tilde{m}_1 - 1); \mathcal{S}) \leq \tau)$ and $L_1 = \left\lfloor \frac{\tilde{T}_1}{\tilde{m}_1 - 1} \right\rfloor$

If \hat{Q}_{t_1} is an estimate of Q_{t_1} with $|\hat{Q}_{t_1} - Q_{t_1}| \leq \beta_{t_1}$ and τ is such that $1 - \hat{Q}_2(\tau) \leq 0.1$ then

$$\left| \mathbb{P}(S_{\tilde{m}_1}(\tilde{T}_1, \mathcal{S}) \leq \tau) - (2\hat{Q}_2 - \hat{Q}_3) \left[1 + \hat{Q}_2 - \hat{Q}_3 + 2(\hat{Q}_2 - \hat{Q}_3)^2 \right]^{1-L_1} \right| \leq E_{total}(1),$$

$$E_{total}(1) = (L_1 - 1) \left[\beta_2 + \beta_3 + F(\hat{Q}_2, L_1 - 1) (1 - \hat{Q}_2 + \beta_2)^2 \right].$$



OUTLINE

1 INTRODUCTION

- Framework
- Problem

2 METHODOLOGY

- Approximation
- Simulation methods: Normal data

3 SIMULATION STUDY

- Numerical examples
- Power

4 SCANNING THE SURFACE OF A CYLINDER

- Problem
- Approximation
- Numerical Results

5 REFERENCES

Numerical examples

SCANNING A REGION OF SIZE $T_1 \times T_2 = 300 \times 350$ TABLE 8 : Numerical results for $\mathbb{P}(S \leq \tau)$: Cylinder**Cylinder:** ($m_1 = 10$, $m_2 = 15$, $T_1 = 300$, $T_2 = 350$, $IS = 1e4$, $IA = 1e5$)

$X_{s_1, s_2} \sim \mathcal{B}(1, 0.1)$				$X_{s_1, s_2} \sim \mathcal{B}(5, 0.05)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal
33	0.871559	0.870200	0.003674	68	0.955593	0.955671	0.000938
34	0.946216	0.946527	0.001177	69	0.976348	0.976285	0.000461
35	0.979458	0.979381	0.000393	70	0.987406	0.987574	0.000227
36	0.992379	0.992465	0.000131	71	0.993526	0.993593	0.000111
37	0.997384	0.997367	0.000043	72	0.996772	0.996751	0.000054
38	0.999116	0.999111	0.000014	73	0.998401	0.998391	0.000026
39	0.999713	0.999714	0.000004	74	0.999214	0.999212	0.000012
40	0.999911	0.999911	0.000001	75	0.999623	0.999626	0.000006

SCANNING A REGION OF SIZE $T_1 \times T_2 = 300 \times 350$ TABLE 8 : Numerical results for $\mathbb{P}(S \leq \tau)$: Cylinder**Cylinder:** ($m_1 = 10, m_2 = 15, T_1 = 300, T_2 = 350, IS = 1e4, IA = 1e5$)

$X_{s_1, s_2} \sim \mathcal{B}(1, 0.1)$				$X_{s_1, s_2} \sim \mathcal{B}(5, 0.05)$			
τ	Sim	AppH	ETotal	τ	Sim	AppH	ETotal
33	0.871559	0.870200	0.003674	68	0.955593	0.955671	0.000938
34	0.946216	0.946527	0.001177	69	0.976348	0.976285	0.000461
35	0.979458	0.979381	0.000393	70	0.987406	0.987574	0.000227
36	0.992379	0.992465	0.000131	71	0.993526	0.993593	0.000111
37	0.997384	0.997367	0.000043	72	0.996772	0.996751	0.000054
38	0.999116	0.999111	0.000014	73	0.998401	0.998391	0.000026
39	0.999713	0.999714	0.000004	74	0.999214	0.999212	0.000012
40	0.999911	0.999911	0.000001	75	0.999623	0.999626	0.000006

SCANNING A REGION OF SIZE $T_1 \times T_2 = 300 \times 350$ TABLE 9 : Numerical results for $\mathbb{P}(S \leq \tau)$: CylinderCylinder: ($m_1 = 10, m_2 = 15, T_1 = 300, T_2 = 350, IS = 1e4, IA = 1e5$)

τ	Sim	AppH	E Total
68	0.915283	0.915691	0.002055
69	0.951447	0.951723	0.001023
70	0.973445	0.973488	0.000515
71	0.985486	0.985509	0.000263
72	0.992349	0.992285	0.000133
73	0.995950	0.995979	0.000066
74	0.997916	0.997920	0.000033
75	0.998951	0.998945	0.000016

SCANNING A REGION OF SIZE $T_1 \times T_2 = 300 \times 350$ TABLE 9 : Numerical results for $\mathbb{P}(S \leq \tau)$: CylinderCylinder: ($m_1 = 10, m_2 = 15, T_1 = 300, T_2 = 350, IS = 1e4, IA = 1e5$)

τ	Sim	$X_{s_1, s_2} \sim \mathcal{P}(0.25)$	ETotal
68	0.915283	0.915691	0.002055
69	0.951447	0.951723	0.001023
70	0.973445	0.973488	0.000515
71	0.985486	0.985509	0.000263
72	0.992349	0.992285	0.000133
73	0.995950	0.995979	0.000066
74	0.997916	0.997920	0.000033
75	0.998951	0.998945	0.000016

thank you!

-  Alm, S. E. (1983).
On the distribution of scan statistic of poisson process.
In Gut, A. and Helst, L., editors, *Probability and Mathematical Statistics*, pages 1–10. Upsalla University Press.
-  Alm, S. E. (1997).
On the distributions of scan statistics of a two-dimensional poisson process.
Advances in Applied Probability, pages 1–18.
-  Alm, S. E. (1998).
Approximation and simulation of the distributions of scan statistics for poisson processes in higher dimensions.
Extremes, 1(1):111–126.
-  Amărioarei, A. (2014).
Approximations for the multidimensional discrete scan statistics.
PhD thesis, University of Lille 1.
-  Anderson, N. H. and Titterington, D. M. (1997).
Some methods for investigating spatial clustering, with epidemiological applications.
Journal of the Royal Statistical Society: Series A (Statistics in Society), 160(1):87–105.

-  Assunção, R., Costa, M., Tavares, A., and Ferreira, S. (2006).
Fast detection of arbitrarily shaped disease clusters.
Statistics in Medicine, 25(5):723–742.
-  Bresenham, J. (1965).
Algorithm for computer control of a digital printer.
IBM Systems Journal, 4(1).
-  Bresenham, J. (1977).
A linear algorithm for incremental digital display of circular arcs.
Communications of the ACM, 20(2):100–106.
-  Devroye, L. (1986).
Non uniform random variate generation.
Springer-Verlag, New York.
-  Foley, J. (1995).
Computer Graphics, Principles and Practice in C.
Addison-Wesley Professional.
-  Glaz, J., Naus, J., and Wallenstein, S. (2001).
Scan Statistics.

Springer.



Glaz, J., Pozdnyakov, V., and Wallenstein, S. (2009).

Scan Statistics: Methods and Applications.
Birkhäuser Boston.



Loader, C. R. (1991).

Large-deviation approximations to the distribution of scan statistics.
Advances in Applied Probability, pages 751–771.



Tango, T. and Takahashi, K. (2005).

A flexibly shaped spatial scan statistic for detecting clusters.

Int J Health Geogr, 4:11.

Tango, Toshiro Takahashi, Kunihiko England Int J Health Geogr. 2005 May 18;4:11.